

**Enhanced Weight Loss from a Dietary Supplement Containing Standardized
Phaseolus vulgaris Extract In Overweight Men and Women**

Running Title: Amylase Inhibitor and Weight Management

Duo Li MSc, PhD¹

Jianguo Shen MD²

Xiaofeng Xu MS³

Alessandra D'Amore MD⁴

Harry G. Preuss, MD, MACN, CNS⁵

1. Department of Food Science and Nutrition
College of Biosystem Engineering and Food Science
Zhejiang Medical University, Hangzhou 310003
2. Department of Endocrinology
The First Affiliated Hospital, College of Medicine
Zhejiang Medical University, Hangzhou 310003
3. Director of Physical Examination Center
Red Cross Hospital
Hangzhou 310003
4. Director of Nuote Nutrition Center
The First Affiliated Hospital, College of Medicine
Zhejiang Medical University, Hangzhou 310003
5. Georgetown University Medical Center
Department of Physiology
Washington, DC 20057

Proofs to:

Harry G. Preuss MD
Professor of Physiology, Medicine, & Pathology
Basic Science Building, Room 231B
Georgetown University Medical Center
Washington, DC 20057

ABSTRACT

Background

One billion human adults worldwide are overweight and, thus, more prone to develop cardiovascular diseases, diabetes, and a variety of other chronic perturbations. Dietary supplements that inhibit gastrointestinal amylase are referred to as "starch blockers" and have been associated with weight loss. Starch blockers may promote weight loss by interfering with breakdown of complex carbohydrates (CHO) limiting their immediate gastrointestinal absorption. Such a process could lessen caloric intake, augment distal CHO fermentation, and/or enhance insulin sensitivity to account for weight loss.

Aims

The present study examines the ability of *Phaseolus vulgaris* extract derived from the white kidney bean to cause weight loss. This extract has previously been shown to inhibit the activity of the digestive enzyme alpha amylase and inhibit starch reabsorption.

Methods

A randomized, double-blinded, placebo-controlled study was conducted in China on 101 volunteers with a BMI between 25-40. The volunteers were divided into two groups that received either placebo or the active substance. A single capsule containing *Phaseolus vulgaris* extract (1,000 mg) or placebo was taken three times per day (just before each meal) for 60 consecutive days. Body weights, and waist and hip measurements were obtained.

Results

After 30 days, 51 subjects receiving *Phaseolus vulgaris* extract compared to a placebo group of 50 subjects had clinically and statistically significantly greater average reduction of body weight [-1.9Kg vs. -0.4Kg, (p<0.001)] and waist circumference [-1.9cm vs. -0.4cm, (p<0.001)], but no difference in the changes of average hip circumference [-0.3cm vs. -0.3cm, (p=0.84)].

Conclusion

The results indicate that *Phaseolus vulgaris* extract taken under the conditions of study produces clinically and statistically significant decrements in body weight and waist circumference.

Key Words:

Starch blockers, weight loss

Obesity, starch blockers

Weight loss, amylase inhibitors

Bean extract, weight loss

INTRODUCTION

It is generally recognized that overweight and obesity have reached epidemic proportions. The proof lies in the fact that there are over one billion overweight adults throughout the world according to many sources including the World Health Organization (WHO) [1-4]. Strategies to lose body fat typically involve a combination of dietary changes limiting caloric intake, physical activity, behavioral therapy, pharmacotherapy, and, in extreme cases, surgery [5]. Since many prefer a natural approach using dietary means to reverse fat accumulation, the availability and popularity of new dietary regimens and natural dietary supplements has risen dramatically in recent years [6,7].

Seeking a safe, easy-to-follow, effective remedy, many have turned to diets proportionately low in refined carbohydrates (CHO) [8-10]. This is based upon continual emergence of data supporting a positive correlation between high refined CHO intake and obesity [11]. Diets possessing stringent depletion of dietary CHO have been tried and shown some success [11]. However, problems arise from such usage: many individuals find such diets unpalatable, while others fear the substitution of too much fat for the missing CHO will increase cardiovascular risk factors. If CHO must remain in the diet for the sake of palatability and the need to lessen fat intake, one alternative is to reduce the gastrointestinal absorption of CHO. Starch blockers are believed to promote weight loss by interfering with and slowing the breakdown of complex CHO, thereby reducing the digestive availability of CHO-derived calories, augmenting distal CHO fermentation, and/or favorably influencing the glucose-insulin system [12-14].

The present paper reports findings from a randomized, single-blinded, placebo-controlled investigation in which body weights of generally healthy, overweight human volunteers in China were examined before and after 30 and 60 days of oral treatment with placebo or a starch blocker produced from white beans¹. While the rationale seems simple, what appears to be a sound hypothesis remains an elusive one to prove. Conclusive, difficult-to-refute results concerning the inhibitory and/or hypoglycemic effects of a bean extract starch blocker are limited and sometimes difficult to interpret [12,13,15-22].

¹ Phase 2 Starch NeutralizerTM, also known as Phaseolamin 2250TM.

Methods

Subject Selection

The present study is randomized, single-blinded, and placebo-controlled investigation carried out with applicable laws related to the protection of study subjects. A total of 101 subjects (50 in the Placebo group and 51 in the Active group) were recruited in Hangzhou, China according to the following inclusion criteria. The healthy subjects, between 20 and 50 years of age, possessed a BMI between 25-40. Subjects were not undergoing any other treatment and had the ability and willingness to understand informed consent and to comply with study procedures. Exclusion criteria included the presence of the following health conditions: diabetes, pregnancy or lactation, active invasive malignant carcinoma, serious hypertension and/or cardiovascular diseases, serious hepatic perturbations, renal damage and severe allergies.

Test Products

The active substance to be tested was a 1,000-mg capsule containing approximately 1,000mg of Phase 2 Starch Neutralizer IV, a dried aqueous extract of the common bean (*Phaseolus vulgaris*). Phase 2 had been standardized to a minimum of 3000 AAIU (alpha-amylase inhibiting units), validated by a modified USP method (SOP 110, Rev.). The control substance was a capsule containing flour containing a similar amount and with the same appearance of the active capsule.

Measurements

Body weight, waist and hip circumferences were measured at the beginning, at 30 days and at the end of the 60-day treatment phase. BMI was calculated taking weight and height into consideration.

Body Weight

Body weights, performed on lightly dressed individuals, were measured using a single, common scale.

Waist and Hip Circumferences

The respective circumference of the waist and hips was measured using a standard non-

stretchable flexible measuring tape. Temporary marks were used to identify the area of reference from one reading to the next.

Adverse Side Effects

The study staff monitored subjects throughout the investigation for the occurrence of any adverse side effects.

Statistical Analysis

At completion, data from 51 subjects receiving the active supplement and 50 subjects receiving the placebo were available for statistical analysis. For each subject, the differences between pre-treatment (baseline) and post-treatment (30 and 60-day) values for each parameter (body weight, waist and hip measurements.) were calculated. The differences were always obtained by subtracting the 30 or 60-day values from the baseline values. A negative difference indicates a reduction in the parameter after 30 or 60 days. Conversely, a positive difference indicates an increase in that parameter. This approach allowed test variability to remain low and the statistical analysis to be more powerful. The differences between pretreatment and 30 and 60-day values were analyzed using Student's t-test. Differences between the Placebo and Active group at each time point were assessed by Student's t test.

Results

The Active and Placebo groups were not statically significantly different in starting weight [79.2 Kg \pm 1.6 (SEM) vs. 81.3 Kg \pm 1.7 (SEM)], height [166.1 cm \pm 1.0 (SEM) vs. 168.1cm \pm 1.1 (SEM)], and age [38.1 years \pm 1.7 (SEM) vs. 39.1 years \pm 1.8 (SEM)] (Tables 1&2). Considering gender, the distribution between Active and Placebo was 27F/24M vs. 19F/31M (Table 1). No significant adverse events were reported. Forty-seven of 51 subjects in the Active group lost weight (92%) after two months, compared to 31 in the Placebo group (62%). The average weight loss in the Active group was 1.9kg compare to a 0.4Kg loss in the Placebo group (p<0.001). There were a lesser percentage of females in the Placebo group (38%) than in the Active group (53%). However, when the weight changes were assessed according to gender, it was found that weight changes were comparable between females and males (Fig. 1).

[Insert Tables 1 &2 and Fig. 1]

The waist size decreased significantly more in the test group (. -1.9 cm vs. -0.4cm, p <0.001), while there were no significant differences in hip size changes between the Test and Placebo groups.

Table 3 gives the blood chemistry values from baseline to one and two months. In general, the chemistries did not change markedly over two months, and there were no statistically significant differences between the Active and the Placebo groups with only two exceptions. Creatinine values at one month were higher in the Active compared to the Placebo group, and the GPT levels averaged lower in the Active than Placebo group. However, these differences were not statistically significant in the two-month values.

[Insert Table 3]

Discussion

Continual emergence of data supporting a positive correlation between excess refined CHO intake and obesity has made many investigators seek more practical means to duplicate results found with the stringent depletion of CHO in the diet [11]. One alternative is use of natural dietary supplements that can considerably lessen starch absorption – thus, mimicking low CHO diets. Ingredients causing such a phenomenon are often referred to as “starch blockers.” We have previously shown in rat and pig models the ability of so-called starch blockers to prevent early absorption of rice starch and sucrose [23,24]. Perhaps the best known of the carb blockers is an extract from bean, [12,13,15-22].

The mechanism behind the weight loss associated with the taking of a starch blocker relies no doubt on alpha-amylase-inhibiting activity [25-29]. *Phaseolus vulgaris* extract has been shown *in vitro* to inhibit the activity of alpha-amylase and may help promote weight loss by interfering with the digestion of complex CHO to simple, absorbable sugars, potentially reducing carbohydrate-derived calories [12,13]. Before crossing the intestinal wall, all complex CHO (*i.e.*, starches) must be hydrolyzed to their mono-

saccharide units, in most cases glucose [30]. There are several enzymes involved in this process: alpha-amylase present in saliva and pancreatic juice, which converts complex CHO into oligosaccharides, and various other enzymes (maltase, lactase, *etc.*) present in the brush border of the small intestine that convert these oligosaccharides to monosaccharides that can then be absorbed.

Unfortunately, data relating carb blockers to weight loss are rare, and those that exist do not always present positive correlations. Earlier studies in the 1980's produced poor results now attributed to insufficient alpha amylase activity in the preparations used at that time [12,13,15-18]. Obviously, the bean extract possessing a greater anti-amylase activity used in starch blocking is very important. In 2007, a dietary formula containing 445 mg of a bean extract taken daily by overweight human subjects concurrently with a carbohydrate-rich, 2000- to 2200-calorie diet was more effective at reducing body weight and body fat mass than placebo [22]. This positive study was carried out with well-studied bean extract named Phase 2. Chokshi noted that Phase 2 is prepared using thermoprocessing conditions to substantially inactivate hemagglutinating activity and trypsin inhibitory activity while preserving substantial alpha-amylase inhibition activity [31,32]. Meiss fed Phase 2 bean extract for 30 days and found a 4% decrease in body weight, accompanied by a 10-45% reduction in body fat [33]. Using the same bean extract, Udani *et al.* [21] reported reduced body weights and serum triglycerides in 14 obese adults receiving 1500 mg of *Phaseolus vulgaris* extract (Phase 2™) twice daily. However, other dose levels were less effective in their study [21].

The present study adds further credibility to the capability of Phase 2 to cause weight loss. After 2 months of receiving one gram of Phase 2 three times a day, the test group lost -1.9 kg vs. -0.4Kg in the placebo group after two months. While baseline values were fairly consistent between the Active and Placebo groups, there were fewer females in the Placebo group. Could this be responsible for the greater weight loss? No! Examining the weight changes separately in each group showed virtually the same changes in each gender as the overall results.

In examining so-called “weight loss studies”, I have repeatedly emphasized that the weight loss is only beneficial if it involves principally fat loss not muscle loss [34]. Although no direct measurements of fat mass were made in the present study, the fact that waist size decreased signifi-

cantly more in the Active group compared to the Placebo group (-1.9cm vs. -0.4cm), whereas, there were no significant differences in hip size suggests that fat loss played a major role in the weight changes.

That this study was performed in China brings up an interesting speculation. The Chinese diet consists largely of rice. Accordingly, the participating subjects were consuming a diet high in CHO. In the successful Celleno study [22], the subjects were supposed to eat a major meal high in CHO content. These associations present a paradox. While we previously point out that a low CHO diet may be best for weight loss [8-10], a diet relatively high in CHO may be preferable when using starch blockers. The latter phenomenon has a number of possible explanations. Perhaps the presence of CHO in the diet is more satisfying to the appetite, replaces other macronutrient calories, and enhances distal fermentation of starches while still performing the functions explained above that create weight loss. A better effect of starch blockers in the presence of a high CHO diet needs further definitive evidence.

In conclusion, use of starch blockers may prove useful in allowing more CHO to be present in the diet and yet obtain results similar to those in diets now restricted in CHO [21,33,35,36].

Acknowledgements.

Drs. Li, Shen, Xu, and D'Amore designed the study and carried it out in China. Dr. Preuss analyzed the data and wrote the manuscript.

Bibliography

1. US Dept of Health and Human Services publication PHS 88-50210, Health objectives for the national prevalence of overweight among adolescents- United States 1988-91, 1994.
2. Popkin BM, Paeratakul S, Zhai F, Keyou G. A review of dietary and environmental correlates of obesity with emphasis on developing countries. *Obes Res* 3: 145S-153S, 1995.
3. WHO Consultation on Obesity. WHO, Geneva, Switzerland, 1997.

4. Guterman L: Obesity problem swells worldwide. *The Chronicle of Higher Education*. A18, March 8, 2002.
5. Bray GA. Obesity. In: Ziegler EE, Filer, Jr LJ Eds, *Present Knowledge in Nutrition*. Washington, D.C., ILSI Press, pp19-32, 1996.
6. Preuss HG, Gottlieb W: *The Natural Fat Loss Pharmacy*. Broadway Books, New York, NY, pp 1-260, 2007 .
7. Agarwal S: Nutritional and Dietary approaches for weight management: an overview. In: *Obesity: Epidemiology, Pathophysiology, and Prevention*. (Eds) D Bagchi, HG Preuss, CRC Press, Fl pp 233-238, 2007.
8. Pawlak DB, Kushner JA, Ludwig DS: Effects of dietary glycaemic index on adiposity, glucose homeostasis, and plasma lipids in animals. *Lancet* 364:778-785, 2004.
9. Brehm BJ, Seeley RJ, Daniels SR, D'Allessio DA: A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *J Fam Pract* 52:515-516, 2003.
10. Meckling KA, Gauthier M, Grubb R, Sanford J: Effects of a hypocaloric, low carbohydrate diet on weight loss, blood lipids, blood pressure, glucose tolerance, and body composition in free-living overweight women. *Canad J Physiol Pharmacol* 80:1095-1105, 2002.
11. Bell SJ, Van Ausdal W, Grachoski G: Appetite, body weight, health implications of a low glycemic load diet. In: *Obesity: Epidemiology, Pathophysiology, and Prevention*. (Eds) D Bagchi, HG Preuss, CRC Press, Fl , pp 233-238, 2007.
12. Layer P, Carlson GL, DiMagno EP: Partially purified white bean amylase inhibitor reduces starch digestion in vitro and inactivates intraduodenal amylase in humans. *Gastroenterology* 88:1895-1902, 1985.
13. Layer P, Zinsmeister AR, DiMagno EP: Effects of decreasing intraluminal amylase activity on starch digestion and postprandial gastrointestinal function in humans. *Gastroenterology* 91:41-48, 1986.
14. Higgins JA, Higbee DR, Donahoo WT, Brown IL, Bell ML, Bessesen DH: Resistant starch consumption promotes lipid oxidation. *Nutrition and Metabolism* 1:8, 2004.

15. Layer P, Rizza RA, Zinsmeister AR, Carlson GL, DiMagno EP: Effect of a purified amylase inhibitor on carbohydrate tolerance in normal subjects and patients with diabetes. *Mayo Clin Proc* 61:442-447, 1986.
16. Boivin M, Zinsmeister AR, Go VI, DiMagno E: Effect of a purified amylase inhibitor on carbohydrate metabolism after a mixed meal in healthy humans. *Mayo Clin Proc* 62:249-255, 1987.
17. Brugge WR, Rosenfeld MS: Impairment of starch absorption by a potent amylase inhibitor. *Am J Gastroenterology* 82:718-722, 1987.
18. Boivin M, Flourie B, Rizza RA, Go VL: Gastrointestinal and metabolic effects of amylase inhibition in diabetics. *Gastroenterology* 94:387-394, 1988.
19. Birketvedt GS, Travis A, Langbakk B, Florholmen JR: Dietary supplementation with bean extract improves lipid profile in over weight and obese subjects. *Nutrition* 18;729-733, 2002.
20. Thom E: A randomized double-blind, placebo-controlled trial of a new weight reducing agent of natural origin. *J Int Med Res* 28:229-233, 2000.
21. Udani J, Hardy M, Madsen DC: Blocking carbohydrate absorption and weight loss: a clinical trial using Phase 2 brand proprietary fractionated white bean extract. *Altern Med Rev* 9:63-69, 2004.
22. Celleno L, Perricone NV, Preuss HG: Effect of a dietary supplement containing standardized *Phaseolus vulgaris* extract on the body composition of overweight men and women. *Int J Med Sci* 4:45-52, 2007.
23. Preuss HG, Echard B, Talpur N, Talpur F, Stohs S: Inhibition of starch and sucrose gastrointestinal absorption in rats by various dietary supplements alone and combined. *Acute studies Int J Med Sci* 4:196-202, 2007.
24. Preuss HG, Echard B, Talpur N, Talpur F, Stohs S: Inhibition of starch and sucrose gastrointestinal absorption in rats by various dietary supplements alone and combined. *Subchronic studies. Int J Med Sci* 4:209-215, 2007.
25. Frels JM, Rupnow JH: Purification and partial characterization of two alpha-amylase inhibitors from black bean (*Phaseolus vulgaris*). *J Food Biochem* 1:385-401, 1984.
26. Gibbs B, Alli I: Characterization of a purified alpha-amylase inhibitor from white kidney bean (*Phaseolus vulgaris*). *Food Research International*. 31:217-225, 1998.

27. Hansawasdi C, Kawabata J, Kasai T: Alpha-amylase inhibitors from roselle (*Hibiscus sabdariffa* Linn.) tea. *Biosci Biotechnol Biochem* 64:1041-1043, 2000.
28. Hansawadi C, Kawabata J, Kasai T: Hibiscus acid as an inhibitor of starch digestion in the Caco-2 cell model system. *Biosci Biotechnol Biochem* 65:2087-2089, 2001.
29. Santimone M, Koukiekolo R, Moreau Y, Le Berre V, Rouge P, Marchis-Mouren G, Desseaux V: Porcine pancreatic alpha-amylase inhibition by the kidney bean (*Phaseolus vulgaris*) inhibitor (Alpha-AII) and structural changes in the alpha-amylase inhibitor complex. *Biochim Biophys Acta* 1696:181-190, 2004.
30. Macdonald I: Carbohydrates. In *Modern Nutrition in Health and Disease*. Edited by Shils ME, Olson JA, Shike M. Pennsylvania: Lea & Febiger; pp 36-46, 1994.
31. Chokshi D: Toxicity studies of Blokal, a dietary supplement containing Phase 2 Starch Neutralizer (Phase 2), a standardized extract of the common white kidney bean (*Phaseolus vulgaris*). *Int J Toxicol* 25:361-371, 2006.
32. Chokshi D: Subchronic oral toxicity of a standardized white kidney bean (*Phaseolus vulgaris*) in rats. *Food Chem Toxicol.* 45:32-40, 2007.
33. Meiss DE: *Phaseolus vulgaris* and alpha-amylase inhibition. In: *Obesity: Epidemiology, Pathophysiology, and Prevention*. (Eds) D Bagchi, HG Preuss, CRC Press, Fl, pp 423-432, 2007.
34. Kaats GR, Michelak J E, Preuss HG: An evaluation of the efficacy and safety of a chitosan fiber product. *J Am Coll Nutr* 25:389-394, 2006.
35. Udani J, Hardy M, Kavoussi B: Dietary supplement carbohydrate inhibitors: a review of the literature. In: *Obesity: Epidemiology, Pathophysiology, and Prevention*. (Eds) D Bagchi, HG Preuss, CRC Press, Fl, pp 279-298, 2007.
36. Obiro WC, Zhang T, Jiang B: The nutraceutical role of the *Phaseolus vulgaris* alpha-amylase inhibitor. *Br J Nutr* 100:1-12, 2008.




Table 1 **Baseline Data at Initiation of Study**

	Active	Placebo
Age (years)	38.1 ± 1.7	39.1 ± 1.8
Gender (F/M)	27/24	19/31
Height (cm)	166.1 ± 1.0	168.1 ± 1.1
BMI	28.6 ± 0.4	28.7 ± 0.4

Ave ± SEM is shown

Table 2 BODY WEIGHT AND MEASUREMENT

	Placebo	Active	p
<u>Scale Weight (Kg)</u>			
Baseline	81.3 ± 1.7	79.2 ± 1.6	0.38
Delta (Month 1 – Baseline)	-0.2 ± 0.12	-1.1 ± 0.10	<0.001
Delta (Month 2 – Baseline)	-0.4 ± 0.13	-1.9 ± 0.15	<0.001
<u>Waist Measurement (cm)</u>			
Baseline	96.9 ± 1.4	95.9 ± 1.5	0.63
Delta (Month 1 – Baseline)	-0.4 ± 0.24	-1.2 ± 0.21	0.01
Delta (Month 2 – Baseline)	-0.4 ± 0.26	-1.9 ± 0.32	<0.001
<u>Hip Measurement (cm)</u>			
Baseline	106.8 ± 1.2	106.9 ± 1.3	0.95
Delta (Month 1 – Baseline)	-0.2 ± 0.15	-0.3 ± 0.23	0.68
Delta (Month 2 – Baseline)	-0.3 ± 0.17	-0.3 ± 0.24	0.84

Average ±SEM is shown

Table 3. **Blood Chemistries at Different Time Points**

Blood Test	Group	Baseline	One Month	Two Months
Glucose (mmol/l)	Pl	5.0 \pm 0.10	4.8 \pm 0.10	4.7 \pm 0.10
	A	4.9 \pm 0.08	4.7 \pm 0.08	4.8 \pm 0.11
	P value	0.53	0.85	0.36
BUN (mmol/l)	Pl	4.5 \pm 0.18	4.6 \pm 0.18	5.4 \pm 0.21
	A	4.6 \pm 0.24	5.2 \pm 0.24	4.9 \pm 0.18
	P value	0.62	0.60	0.60
Creatinine (umol/l)	Pl	70.3 \pm 2.5	74.7 \pm 3.3	82.2 \pm 2.2
	A	75.5 \pm 2.5	86.4 \pm 2.7	85.5 \pm 2.7
	P value	0.15	0.01	0.34
Uric Acid (mmol/l)	Pl	218 \pm 10.9	248 \pm 9.8	226 \pm 10.1
	A	239 \pm 10.5	236 \pm 8.8	225 \pm 9.0
	P value	0.17	0.36	0.94
T Chol (mmol/l)	Pl	4.9 \pm 0.09	4.6 \pm 0.09	4.7 \pm 0.09
	A	4.9 \pm 0.08	4.6 \pm 0.08	4.7 \pm 0.08
	P value	0.88	0.84	0.53
LDL (mmol/l)	Pl	3.4 \pm 0.09	3.4 \pm 0.07	3.4 \pm 0.07
	A	3.4 \pm 0.10	3.3 \pm 0.07	3.4 \pm 0.08
	P value	0.76	0.44	0.99
HDL (mmol/l)	Pl	1.4 \pm 0.04	1.4 \pm 0.04	1.4 \pm 0.04
	A	1.5 \pm 0.06	1.5 \pm 0.04	1.5 \pm 0.04
	P value	0.49	0.49	0.13

Trigly (mmol/l)	Pl	1.7±0.11	1.5±0.09	1.5±0.09
	A	1.7±0.18	1.5±0.13	1.5±0.10
	P value	0.95	0.29	0.34
Apo A (mmol/l)	Pl	1.4±0.03	1.4±0.03	1.4±0.03
	A	1.4±0.03	1.4±0.03	1.4±0.03
	P value	0.57	0.41	0.78
Apo B (mmol/l)	Pl	1.2±0.03	1.2±0.03	1.2±0.03
	A	1.2±0.03	1.2±0.03	1.2±0.03
	P value	0.75	0.71	0.19
GPT (U/l)	Pl	42.5±2.5	40.7±1.5	35.8±1.8
	A	41.1±1.9	35.3±1.4	32.4±1.3
	P value	0.66	0.01	0.12
AAT (U/l)	Pl	41.4±1.8	37.4±1.5	32.4±1.4
	A	42.9±1.6	37.8±1.5	31.9±1.4
	P value	0.52	0.86	0.79
GIT (U/l)	Pl	37.5±2.5	35.2±3.3	30.6±2.2
	A	38.8±2.5	39.0±2.7	32.8±2.7
	P value	0.61	0.08	0.19

Values are in SI units. Average ± SEM is shown. P value is for difference between placebo and active at indicated time point. BUN=blood urea nitrogen, T Chol= total cholesterol, Trigly=triglycerides, Apo A – apolipoprotein A, Apo B= apolipoprotein B, GPT=glutamate pyruvate transaminase, AAT=aspartate aminotransferase, GIT= glutamyltranspeptidase